

Virulence profiling of *Campylobacter* spp *Campylobacter jejuni* and *C. fetus* subsp *fetus*. abortions rise in sheep farms in Kashmir, India

M. Hafiz¹, S. Qureshi*¹, M. Gulzar¹, Z. Kashoo¹, M. Sharief Bandy², S. Farooq¹,
M. Altaf Bhat¹, P. Dar¹, S.A. Hussain³, S.M. Andrabi⁴, M.I. Hussain¹, G. Badroo¹, F. ud Din¹

¹ Campylobacter Laboratory; Division of Veterinary Microbiology and Immunology, Faculty of Veterinary Sciences and Animal Husbandry, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir (SKUAST-K), Shuhama (Aulesteng)-19006, Jammu and Kashmir, India

² Department of Clinical Pharmacology, Sher-i-Kashmir Institute of Medical Sciences, Soura, Srinagar- 190011, Jammu and Kashmir, India

³ Division of Veterinary Public Health and Epidemiology, Faculty of Veterinary Sciences and Animal Husbandry, Sher-e-Kashmir University of Agricultural Sciences & Technology of Kashmir (SKUAST-K) Shuhama (Aulesteng) Srinagar-19006, Jammu and Kashmir, India

⁴ Division of Animal Biotechnology, Faculty of Veterinary Sciences and Animal Husbandry, Sher-e -Kashmir University of Agricultural Sciences and Technology of Kashmir (SKUAST-K) Shuhama (Aulesteng) Srinagar-19006, Jammu and Kashmir, India

Correspondence to: S. Qureshi, e-mail: qureshi.sabia@gmail.com

Abstract

Campylobacter spp. are the leading causes of ovine abortions leading to severe economic losses and a source of bacterial food borne illness in humans, posing a major public health concern. This study reports an increase in *Brucella* negative abortions in sheep farms in Kashmir, India in the last few years. Screening of sheep farms was carried to rule out *Campylobacter* abortion. Three Government sheep breeding farms in the Kashmir valley and some other private flocks were screened for the presence of *C. jejuni* and *C. fetus* subsp. *fetus*. A total of 217 samples comprising of 200 swabs (rectal and vaginal swabs; 100 each) from clinically healthy animals, ten vaginal swabs from cases of abortion and seven abomasal contents of the aborted fetuses were collected from sheep breeding farm Khimber (District, Srinagar), sheep breeding farm Goabal, the Mountain Research Centre for sheep and goats (MRCSG, SKUAST-K) (District, Ganderbal) and from private sheep breeders were screened. In the present study a total of 15.2% of samples were positive for different *Campylobacter* spp. by PCR. *C. jejuni* and *C. coli* were detected individually or coexisting by PCR in the rectal swabs from all farms, while *C. fetus* subsp. *fetus* was detected only in the vaginal swabs from private sheep farms and abortion samples. Additionally, *C. jejuni* was also isolated from vaginal swabs. A total of 5, 20 and 18 samples were confirmed positive by PCR for *C. fetus* subsp. *fetus*, *C. jejuni* and *C. coli*, respectively. The *Campylobacter* isolates obtained in the present study were screened for *flaA*, *cdtB*, *cadF*, *wlaN*, *pldA*, *virB* and *dnaJ* virulence determinants. However, the isolates harboured *flaA*, *cdtB* and *cadF* virulence determinants only. The recovery of virulent *Campylobacter* isolates from healthy sheep fecal swabs in the present study may have longer human health implications. The presence of abortive strains of *C. jejuni* and *C. fetus* subsp. *fetus* in sheep farms has long term economic implications in the Kashmir valley. This study emphasizes the need for efforts to be taken on farms to prevent animal infections and minimizing human exposure to these pathogens through proper hygiene and production practices as suggested by OIE.

Keywords: Abortion, *C. jejuni*, *C. coli*, *C. fetus* subsp. *fetus*, sheep, virulence determinants



Introduction

The genus *Campylobacter* identified in 1886 by Theodore Escherich was recognized as an etiological agent of abortion in sheep nearly a century ago (McFaydean and Stockman, 1913). *Campylobacter* a gram negative, oxidase, catalase positive and non-fermentative bacterium of the family *Campylobacteraceae* requires microaerophilic atmosphere of 3-5% oxygen, 10% CO₂, and 85% nitrogen for optimal growth, with some species requiring hydrogen (Bolton and Coates, 1983). The recent years have witnessed increased human cases of *Campylobacter* gastroenteritis, making it the second most serious food-borne illness globally after Salmonellosis. *Campylobacter jejuni* and *Campylobacter coli* are commonly found in the intestinal contents or feces of healthy sheep (Mehmut et al. 2006). Although poultry are the largest reservoir for human campylobacteriosis, research has confirmed animal faeces, mutton and milk as a source of human campylobacteriosis for their handlers.

Campylobacter spp. (*C. jejuni* and *C. fetus* subsp. *fetus*) are the primary cause of ovine abortion globally, with an average occurrence rate of 23.2% within affected flocks (Sahin et al. 2008, Joens et al. 2010). *Campylobacter fetus* subsp. *fetus* has been recognized as the leading cause of sheep abortion (Sahin et al. 2017) and *C. jejuni* has also been implicated in sheep abortion (Yaeger et al. 2021).

Several virulence-associated genes play an important role in *Campylobacter* pathogenicity (Zilbauer et al. 2008). The *fla* genes (*flaA* and *flaB*) responsible for bacterial motility, encode flagellin, the ciliary protein which enables *Campylobacter spp.* cells to move and colonize. The *cadF* conserved gene, encoding the fibronectin binding protein of enterocytes, participates in the adherence necessary to induce symptoms of Campylobacteriosis, in *C. jejuni* and *C. coli* (Ziprin et al. 2001). The *vir* gene, encoded by the *Campylobacter* plasmid, encodes proteins responsible for pathogenicity (Bacon et al. 2000). The *pldA* gene having a role in cell invasion is responsible for the synthesis of an outer membrane phospholipase important for caecal colonization. The *dnaJ* considered to be a chaperone protein helps to cope with diverse physiological stresses (Chansiripornchai and Sasipreeyajan 2009). The *ciaB*, *pldA* and *dnaJ* genes, recognised as heat shock protein genes are important for caecal colonisation and mutations in these genes limit the ability of the organism to colonise (Reddy et al. 2018). The cytotoxic distending toxin (*cdt*) genes, *cdtA*, *cdtB* and *cdtC*, form the polycistronic *cdt* operons that are responsible for the expression of cytotoxins and are lethal for host enterocytes. The strains of *C. jejuni* may carry *wlaN* which is responsible for causing Guillain Barre syn-

drome due to its ability to produce LOS (lipooligosaccharide) which exhibits molecular mimicry with the saccharide component of human GMI ganglioside present in peripheral nerves (Perez-Perez et al. 1996, Yuki et al. 2001, Poropatich et al. 2010).

Not much recognition was previously given to this organism as the causative agent of abortion however, it has been found to be the predominant isolate responsible for abortion (Kirkbridge 1993) in sheep in many studies. Sheep rearing is an important source of livelihood especially among migratory Gujjar Bakerwal tribes in the union territory (UT) of Jammu and Kashmir. The consumption of mutton in the Kashmir valley is high due to the severe cold climatic conditions in winter which necessitates the need for local sheep rearing and purchase from other states of the country. The losses due to *Campylobacter* abortion in sheep flocks have not been assessed previously in the UT of J&K. The past decade has witnessed a steady increase in *Brucella* negative abortions in sheep flock in the UT. Several reports of abortions from Government sheep farms in the UT with negative *Brucella* serology of the aborted animals form the basis of this study.

Materials and Methods

Sampling

A total of 200 samples (comprising 25 rectal and 25 vaginal swabs) were collected from sheep breeding farm Khimber, Mountain Research Centre for sheep and goats, sheep breeding farm Goabal, and private sheep breeding farms in the Ganderbal district. Abomasal contents (n=7) from aborted sheep foetus and vaginal swabs from ten abortion cases, which were previously confirmed as *Brucella* negative were also screened in the present study. The samples were collected in sterile vials, and transported to the laboratory on ice in Cary Blairs transport media (Hi Media, India).

Bacterial DNA Extraction and PCR assays

The samples were enriched by inoculating in thio-glycollate broth (Difco) and incubated under microaerophilic conditions using a campygen gas pack (BD Gas Pak Campy Container system) initially at 30°C for 5 hours followed by incubation at 42°C for 24-48 hrs for *C. jejuni/C. coli* and 37°C for 24-48 hrs for *C. fetus* subsp. *fetus*. Enriched samples were streaked on modified Cefaperazone Charcoal Agar (mCCDA) supplemented with mCCDA supplement (Hi Media, India) and plates were incubated under microaerophilic conditions using the temperature and time combination as described previously (Gharbi et al. 2018.). The identification of colonies was based on colony morphology,

Table. 1 Primer sequence used for amplification of 16SrRNA *Campylobacter* genus, *Campylobacter jejuni* and *Campylobacter coli*.

S.No.	Target gene	Primer sequence (5'-3')	Product size (bp)	Reference
1.	16SrRNA (Genus specific)	F-GGTAAAGTCCCGCAACGAGCCGC R-GGCTGATCTACGATTACTAGCGAT	283	Leite et al. (1996)
2.	<i>mapA</i> (<i>C.jejuni</i>)	F-CTATTTTATTTTGGAGTGCTTGTG R-GCTTTATTTGCCATTTGTTTTATTA	589	Denis et al. (1999)
3.	<i>ceuE</i> (<i>C. coli</i>)	F-AATTGAAAATTGCTCCAACATATG R- TGATTTTATTATTTGTAGCAGCG	462	Denis et al. (1999)
4.	<i>CFE</i> (<i>C. fetus</i> subsp. <i>fetus</i>)	F-GCAAATATAAATGTAAGCGGAGAG R-TGCAGCGGCCACCTAT	435	Wang et al. (2002)

Table. 2. Primer sequence for detection of virulence genes of *C. jejuni* and *C. coli*.

S.No.	Target gene	Primer sequence (5'-3')	Product size (bp)	Reference
1.	<i>cdtB</i>	F-CAGAAAGCAAATGGAGTGTT R- AGCTAAAAGCGGTGGAGTAT	620	Hickey et al. (2000)
2.	<i>cadF</i>	F-TTGAAGGTAATTTAGATATG R-CTAATACCTAAAGTTGAAAC	400	Konkel et al. (1999)
3.	<i>flaA</i>	F-AATAAAAATGCTGATAAACAGGTG R-TACCGAACCAATGCTGCTCTGATT	855	Konkel et al. (1999)
4.	<i>dnaJ</i>	F-AAGGCTTTGGCTCATC R-CTTTTTGTTTCATCGTT	720	Suvamoy et al. (2003)
5.	<i>pldA</i>	F-AAGCTTATGCGTTTTT R-TATAAGGCTTTCTCCA	913	Suvamoy et al. (2003)
6.	<i>virB</i>	F-TCTTGTGAGTTGCCTTACCCCTTTT R-CCTGCGTGTCTGTGTTATTTACCC	494	Suvamoy et al. (2003)

gram staining and biochemical tests, (catalase, oxidase and hippurate hydrolysis for *C. jejuni*).

DNA extraction was performed using the phenol chloroform isoamyl (PCI) extraction method. Briefly, suspected colonies was suspended in a 1.5 ml microcentrifuge tube containing 200 µl of distilled water and gently mixed by vortex. The samples were boiled for 5 min, followed by cooling on ice for 10 min followed by centrifugation at 10,000 × g in a table-top microcentrifuge (Cooling Centrifuge, Eppendorf) for 10 min. The supernatants were discarded before adding 250 µl RNase (100 µg/ml) and 250 µl of lysis buffer. A total of 550 µl of saturated phenol was added mixed thoroughly and followed by centrifugation at 8,000 × g. The supernatant was collected and extracted with phenol, chloroform and isoamyl alcohol (25:24:1) and centrifuged at 8,000 × g for 10 min. The supernatant was extracted twice using PCI as above. The aqueous phase was collected followed by cold salt precipitation with 2 M sodium acetate and 1.5 ml ethanol (100%) and kept at -20°C for 1 hr followed by centrifugation at 12,000 × g. The DNA pellet was washed with 80% ethanol, dried and suspended in 30 µl of 1X TAE buffer and used as the template (40-100 ng/µl) for PCR.

The PCR protocol as per Wang et al. (2002) was followed for detection of *C. fetus* subsp. *fetus* while the protocol of Denis et al. (1999) was followed for *Campylobacter jejuni* and *Campylobacter coli* detection. Details of the primers used in this study are provided in Table 1. The cyclic condition for the *sapB* gene amplification of *C. fetus* subsp. *fetus*, involved an initial denaturation at 94°C for 5 minutes, followed by 30 cycles of denaturation at 94°C for 1 minute, annealing at 58°C for 60 seconds, and extension at 72°C for 60 seconds and a final extension step at 72°C for 5 minutes (Wang et al. 2002).

The cyclic conditions for multiplex PCR for *Campylobacter jejuni* and *Campylobacter coli* amplification included an initial denaturation at 94°C for 5 minutes, followed by 35 cycles of denaturation at 94°C for 1 minute, annealing at 52°C for 1 minute, and extension at 72°C for 1 minute and a final extension step at 72°C for 10 minutes (Denis et al. 1999). The *Campylobacter jejuni* and *Campylobacter coli* isolates were screened for virulence genes, *cadF*, *cdtB*, *flaA*, *dnaJ*, *pldA*, and *virB*, using primers as described in Table 2.




Fig. 1. *Campylobacter jejuni* on mCCDA (Modified Charcoal Cefoperazone-deoxycholate) agar.




Fig. 2. Multiplex PCR for *mapA* (*C. jejuni*) and *ceuE* (*C. coli*) genes.

M: 100bp ladder, P1: Positive control *C. jejuni*, P2: Positive control *C. coli*. N: Negative control, 1-2: isolates positive for *Campylobacter jejuni* 5-8: isolates positive for *C. coli*

Antimicrobial sensitivity profile

The antimicrobial sensitivity profile was determined using the standard disc diffusion method on Muller-Hinton agar containing 5% defibrinated sheep's blood. The panel of antibiotic discs included penicillin, amikacin, tetracycline, ciprofloxacin, cephalothin, nalidixic acid and enrofloxacin. The interpretation of the test was done based on M-45; Clinical and Laboratory Standards Institute (Beilei-Ge et al. 2013).

Results

In the present study, out of 33 PCR positive samples only 12 isolates of *Campylobacter spp* could be obtai-

ned. These comprised 7 isolates of *C. jejuni*, 3 isolates of *C. coli*, and 2 isolates of *C. fetus* subsp. *fetus*. Typical small, round, smooth, glistening buff coloured colonies were observed on mCCDA medium (Fig. 1) which revealed Gram negative rods with typical seagull morphology.

A total of 33 (15.2%) out of 217 samples screened from various sheep farms (both organized and unorganized) were found positive for *Campylobacter spp* by PCR. The amplified products of 283bp, 589bp, 462bp, and 435bp corresponding to the genus *Campylobacter*, *C. jejuni*, *C. coli*, and *C. fetus* subsp. *fetus* respectively, were obtained in the PCR assays (Figs. 2, 3).

From sheep breeding farm Goabal, 1 (4%) of the 25 rectal swabs, tested positive for *C. jejuni*, 1 (4%) for




Fig. 3. Agarose gel electrophoresis showing 435bp amplicon of *Campylobacter fetus* subsp. *fetus*.
M: 100bp ladder, P: Positive control, N: Negative control, 1-3: Isolates positive for *Campylobacter fetus* subsp. *fetus*




Fig. 4. Multiplex PCR for *fla* and *Cdt* gene of *Campylobacter* spp.
M: 100bp ladder, P: Positive control *Fla* (855bp) and *Cdt* (620bp) genes, 1-4: *Fla* and *Cdt* gene positive isolates, N: Negative control

C. coli, and 1(4%) sample was positive for both *C. jejuni* and *C. coli*. Of the 25 vaginal swabs collected from the same farm, 1 (4%) tested positive for *C. jejuni* by PCR. From the samples collected at sheep breeding farm Khimber, 3 (12%) of the 25 rectal swabs, were positive for *C. jejuni*, 1 (4%) for *C. coli*, and 2 (8%) has coexistent *C. jejuni* and *C. coli*. None of the 25 vaginal swabs from the farm tested positive for any of *Campylobacter* spp by PCR. In MRCG Shuhama, out of the 25 rectal swabs 3(12%) were positive for *C. jejuni*, 3(12%) for *C. coli*, and 4(16%) for both *C. jejuni* and *C. coli*. None of the 25 vaginal swabs screened from the farm revealed the presence of *Campylobacter* spp. In private sheep farms, out of the 25 rectal swabs 2(8%) tested positive for *C. jejuni*, 3 (12%) for *C. coli*, and 3 (12%) for both *C. jejuni* and *C. coli*. Out of the 25 vaginal swabs screened from these private farms 1(4%) was positive for *C. fetus* subsp. *fetus*. Out of the 7 abo-

masal contents from aborted sheep fetuses, 1(14.2%) tested positive for *C. fetus* subsp. *fetus*. Of the 10 vaginal swabs from *Brucella* negative aborted sheep 3 (30%) were positive for *C. fetus* subsp. *fetus*.

All the isolates of the present study revealed the presence of *cadF*, *flaA* and *cdtB* (Figs. 4, 5). However, none of the isolates harboured the *pldA*, *dnaJ*, *virB*, and *wlaN* gene. The distribution of *Campylobacter* spp in various organized and private farms is shown in Table 3.

The antibiotic sensitivity profile revealed that all isolates were resistant to penicillin (100%), cephalothin (100%), with resistance to tetracycline being 80%, amikacin (76%), ciprofloxacin (68%) and enrofloxacin (52%). In the present study 80% of isolates revealed sensitivity to nalidixic acid.




Fig. 5. Agarose gel electrophoresis showing 400bp amplicon of *CadF* gene.

M: 100bp Ladder; P: Positive control, N: Negative control; 1-4: *CadF* positive samples

Table 3. Distribution of *Campylobacter* spp in various farms.

S.No	Farm name & District	Sample type and no	<i>C. jejuni</i>	<i>C. coli</i>	<i>C. jejuni</i> & <i>C. coli</i>	<i>C. fetus</i> subsp <i>fetus</i>	No. of Isolates		
							<i>C. jejuni</i>	<i>C. coli</i>	<i>C. fetus</i> subsp <i>fetus</i>
1.	Sheep Breeding Farm Goabal, (District Ganderbal)	Rectal swabs (n=25)	1(4%)	1(4%)	1(4%)	0	1	0	0
		Vaginal swabs (n=25)	1(4%)	0	0	0	0	0	0
2.	Sheep Breeding Farm Khimber, (District Srinagar)	Rectal swabs (n=25)	3(12%)	1(4%)	2(8%)	0	2	0	0
		Vaginal swabs (n=25)	0	0	0	0	0	0	0
3.	MRCG Shuhama (District Ganderbal)	Rectal swabs (n=25)	3(12%)	3(12%)	4(16%)	0	1	2	0
		Vaginal swabs (n=25)	0	0	0	0	0	0	0
4.	Private sheep farms	Rectal swabs (n=25)	2(8%)	3(12%)	3(12%)	0	3	1	0
		Vaginal swabs (n=25)	0	0	0	1(4%)	0	0	1
5.	Cases from veterinary hospitals	Abomasal content of the aborted sheep foetus (n=7)	0	0	0	1(14.2%)	0	0	0
6.	Cases from veterinary hospitals	Vaginal swabs from <i>Brucella</i> negative aborted sheep (n=10)	0	0	0	3(30%)	0	0	1

Discussion

Campylobacter species pose a significant threat to both animals and humans, particularly in sheep-rearing regions worldwide. *Campylobacter fetus* subsp. *fetus*

and *C. jejuni* are a major cause of sheep abortion and can lead to severe economic losses in the sheep industry (Hedstrom et al. 1987, Dorsch et al. 2022). Infection occurs through ingestion of the bacteria, with most abortions occurring during the last month of pregnancy.

The genus *Campylobacter* is also recognized as an important zoonotic pathogen, with *C. jejuni* and *C. coli* being a common cause of human gastroenteritis, surpassing cases of salmonellosis in high-income countries (Cobo-Diaz et al. 2021). The disease spreads rapidly among uninfected ewes unless stringent hygiene practices are followed, as the fetus, placenta, birth fluids, vaginal discharge, and feces from the ewe are all potential sources of infection. It has been observed that if water or feeding areas become contaminated with these materials, the abortion rate can be notably high (Agerholm et al. 2006, Mearns et al. 2007).

In the present study the occurrence of 15.2% of *Campylobacter* (33/217) from both organized and private sheep farmers in the Kashmir valley was observed. A high percentage of *C. fetus* subsp. *fetus* was detected in the abortion samples (14.2% from abomasal contents of aborted sheep fetus and 30% from vaginal swabs of aborted sheep) in comparison to the vaginal swabs from healthy animals (1%). *C. jejuni* (1%) was only isolated from vaginal swabs of the healthy animals. Several studies (Bisma et al. 2018 Hamali et al. 2014) have reported the presence of *C. jejuni* and *C. fetus* subsp. *fetus* from the vaginal swabs and aborted samples. Bisma et al. (2018), reported 10% of cases positive for *Campylobacter jejuni*, and 3.33% of cases for *Campylobacter fetus* subsp. *fetus* from 150 vaginal samples of aborted ewes. Hamal et al. (2014) reported a prevalence of 9.09% of cases of *C. jejuni* and 1.51% of cases of *C. fetus* subsp. *fetus* from 132 aborted fetus and placental samples. Out of the 100 rectal swab samples screened from sheep in the same study 19% were *C. jejuni* and 18% were *C. coli*. Adesiyun et al. (1992) reported a prevalence rate of 17.92% *Campylobacter* sp from lamb rectal swabs.

A passive filtration technique using 0.45µ pore size filters post enrichment with campy thioglycollate broth was used for isolation of *Campylobacter*. Only 23.2% (10/33) of *Campylobacter* spp could be isolated from the samples. A plausible reason for this low isolation rate is attributed to its extremely slow growth and fastidious nature (Brand et al. 2004, Chon et al. 2022). All the *C. jejuni* isolates were hippurate hydrolysis positive, and all *C. fetus* subsp. *fetus* isolates were positive in the nitrate reduction test. Harvey et al. (1980) and Steinhauserova et al. (2001) have reported hippurate hydrolysis and nitrate reduction tests as one of the important biochemical tests for identification of *C. jejuni* and *C. fetus* subsp. *fetus*, respectively.

All the isolates of the present study revealed the presence of *fla*, *cad*, *cdt* virulence genes. The pathogenesis of *Campylobacter* spp. particularly *Campylobacter jejuni* and *Campylobacter coli* has been attributed to the presence of several virulence genes viz;

fla, *cad*, *virB11*, *cdt*, *wlaN*, *dnaJ* and *pldA* (Gilbert et al. 2000). The polar flagellum encoded by (*fla*), responsible for the characteristic darting motility of *C. jejuni* is involved in the colonization and invasion (Guerry, 1997), *cdt* which blocks the G2/M phase of eukaryotic cells prior to cell division, induces a cytoplasmic distention and ultimately causes cell death (Jeon et al. 2005), and *Campylobacter* adhesion to fibronectin protein (*cadF*) promotes intestinal epithelial cell binding (Monteville et al. 2003). These virulence determinants in *Campylobacter* strains isolated from various sources has been reported by Bang et al. (2003), Datta et al. (2003) and Muller et al. (2006) emphasizing the important role of these virulence markers in *Campylobacter* pathogenesis.

None of the isolates in the present study revealed the presence of *dnaJ*, *virB11*, *pldA* and *wlaN* genes. Similar findings have been reported by Datta et al. (2002) who analysed 40 *Campylobacter* isolates from different parts of Japan by PCR. All the isolates were found positive for the *fla* gene, 16 (32%) for *cad*, 17 (42.5%) for *racR* and 7 (17.5%) for *cdt*. No samples were positive for *wlaN*. Bisma et al. (2018) also reported the presence of the virulence genes *fla*, *cdt*, and *cad* in seventeen isolates of *C. jejuni* and three isolates of *C. coli*.

Antimicrobial resistance is a major public health concern in both developed and developing countries in recent years (Padungton and Kaneene 2003). The most common antibiotics used for the treatment of *Campylobacter* associated enteritis and abortions in animals are macrolides, fluoroquinolones and tetracycline. A high degree of resistance has been reported in recent years to these antibiotics around the globe (Wieczorek and Osek, 2013). The *in vitro* antibiotic sensitivity profile of *Campylobacter* isolates in the present study revealed highest sensitivity of the isolates to nalidixic acid and resistance to penicillin, cephalothin, tetracycline, amikacin, ciprofloxacin and enrofloxacin. Several researchers such as Bisma et al. (2018), Abdullah et al. (2022) and Chepkwony (2016) have reported the multidrug resistance of *Campylobacter* spp to various classes of antibiotics.

This study emphasizes the need for monitoring and controlling *Campylobacter* infections in sheep farms of the union territory of J&K, India to safeguard animal health and mitigate the risk of zoonotic transmission to humans. Proper management practices and hygienic measures are crucial in curbing the spread of this disease and ensuring the well-being of both the sheep population and humans.

Acknowledgements

We would like to acknowledge the help provided by Prof. Linda Van der Graaf, Department of Infectious Disease & Immunology, Faculty of Veterinary Medicine, Utrecht University, Netherland, for providing the control DNAs. The study was financed by DBT, GOI, Grant No: BT/PR40347/ADV/90/286/2020, dated 22/09/2021.

References

- Abdallah M, Abaza MA, Fathy RR, Youseef AG, Sobhy M, Abd Elhamid HS, Ahmed W (2022) Detection of Some Virulence and Antibiotic Resistance Genes in *Campylobacter jejuni* isolated from Poultry and Human. *Egypt J Hosp Med* 1: 6373-6381.
- Acik MN, Cetinkaya B (2006) Heterogeneity of *Campylobacter jejuni* and *Campylobacter coli* strains from healthy sheep. *Vet Microbiol* 115: 370-375.
- Adesiyun AA, Kaminjolo JS, Loregnard R, Kitson-Piggott W (1992) *Campylobacter* infections in calves, piglets, lambs and kids in Trinidad. *Br Vet J* 148: 547-556.
- Agerholm JS, Aalback B, Fog-Larsen AM, Boye M, Holm E, Jensen TK, Lindhardt T, Larsen LE, Buxton D (2006) Veterinary and medical aspects of abortion in Danish sheep. *APMIS* 114: 146-152.
- Bacon DJ, Alm RA, Burr DH, Hu L, Kopecko DJ, Ewing CP, Trust TJ, Guerry P (2000) Involvement of a plasmid in virulence of *Campylobacter jejuni* 81-176. *Infect Immun* 68: 4384-4390.
- Bang DD, Nielsen EM, Scheutz F, Pedersen K, Handberg K, Madsen M (2003) PCR detection of seven virulence and toxin genes of *Campylobacter jejuni* and *Campylobacter coli* isolates from Danish pigs and cattle and cytolethal distending toxin production of the isolates. *J Appl Microbiol* 94: 1003-1014.
- Bisma G, Sabia Q, Zahid AK, Ahmad WS, Nabi MS, Ahmad MF, Isfaul HM, Aasim H, Ali RM, Shafkat MK, Rafia M (2018) RFLP analysis of flagellin (*Fla*) gene of *Campylobacter jejuni* from ovines of Kashmir, India. *J Food Saf* 38: 12509.
- Bolton FJ, Coates D (1983) A study of the oxygen and carbon dioxide requirements of thermophilic *campylobacters*. *J Clin Pathol* 36: 829-834.
- Brandl MT, Haxo AF, Bates AH, Mandrell RE (2004) Comparison of survival of *Campylobacter jejuni* in the phyllosphere with that in the rhizosphere of spinach and radish plants. *Appl Environ Microbiol* 70: 1182-1189.
- Cardarelli-Leite P, Blom K, Patton CM, Nicholson MA, Steigerwalt AG, Hunter SB, Brenner DJ, Barrett TJ, Swaminathan B (1996) Rapid identification of *Campylobacter* species by restriction fragment length polymorphism analysis of a PCR-amplified fragment of the gene coding for 16S rRNA. *J Clin Microbiol* 34: 62-67.
- Chansiripornchai N, Sasipreeyajan, J (2009) PCR detection of four virulence-associated genes of *Campylobacter jejuni* isolates from Thai broilers and their abilities of adhesion to and invasion of INT-407 cells. *J Vet Med Sci* 71: 839-844.
- Chepkwony MC (2016) Prevalence and Antimicrobial resistance of zoonotic campylobacter isolated from livestock and rodents in urban informal settlements in Nairobi (Doctoral Dissertation, University of Nairobi).
- Chon JW, Hyeon JY, Yim JH, Kim JH, Song KY, Seo KH (2012) Improvement of modified charcoal-cefoperazone-deoxycholate agar by supplementation with a high concentration of polymyxin B for detection of *Campylobacter jejuni* and *C. coli* in chicken carcass rinses. *Appl Environ Microbiol* 78: 1624-1626.
- Chon JW, Seo KH, Kim B, Jeong D, Song KY (2020) Advanced methods for isolating from and confirming *Campylobacter* spp. in milk and dairy products. *J Dairy Sci Biotechnol* 38: 121-133.
- Cobo-Diaz JF, Gonzalez del Rio P, Alvarez-Ordóñez A (2021) Whole resistome analysis in *Campylobacter jejuni* and *C. coli* genomes available in public repositories. *Front Microbiol* 12: 662144.
- Datta S, Niwa H, Itoh K (2003) Prevalence of 11 pathogenic genes of *Campylobacter jejuni* by PCR in strains isolated from humans, poultry meat and broiler and bovine faeces. *J Med Microbiol* 52: 345-348.
- Datta S, Niwa H, Itoh K (2003) Prevalence of 11 pathogenic genes of *Campylobacter jejuni* by PCR in strains isolated from humans, poultry meat and broiler and bovine faeces. *J Med Microbiol* 52: 345-348.
- Denis M, Soumet C, Rivoal K, Ermel G, Blivet D, Salvat G, Colin P (1999) Development of am-PCR assay for simultaneous identification of *Campylobacter jejuni* and *C. coli*. *Lett Appl Microbiol* 29: 406-410.
- Dorsch MA, Casaux ML, Calleros L, Araoz V, Caffarena RD, Monesiglio C, Barcellos M, da Silva Silveira C, Perdomo Y, Banchero G, Uzal FA (2022) Placentitis and abortion caused by a multidrug resistant strain of *Campylobacter fetus* subspecies *fetus* in a sheep in Uruguay. *Rev Argent Microbiol* 54: 25-30.
- Findik A, Ica T, Onuk EE, Percin D, Kevenk TO, Ciftci A (2011) Molecular typing and *cdt* genes prevalence of *Campylobacter jejuni* isolates from various sources. *Trop Anim Health Prod* 43: 711-719.
- Ge B, Wang F, Sjolund-Karlsson M, McDermott PF (2013) Antimicrobial resistance in *Campylobacter*: susceptibility testing methods and resistance trends. *J Microbiol Methods* 95: 57-67.
- Gharbi M, Bejaoui A, Ben Hamda C, Jouin A, Ghedira K, Zrelli C, Hamrouni S, Aouadhi C, Bessoussa G, Ghram A, Maaroufi A (2018) Prevalence and antibiotic resistance patterns of *Campylobacter* spp. isolated from broiler chickens in the North of Tunisia. *Biomed Res Int* 2018: 1-7. <https://doi.org/10.1155/2018/7943786>.
- Gilbert M, Brisson JR, Karwaski MF, Michniewicz J, Cunningham AM, Wu Y, Young NM, Wakarchuk WW (2000) Biosynthesis of ganglioside mimics in *Campylobacter jejuni* OH4384: identification of the glycosyltransferase genes, enzymatic synthesis of model compounds, and characterization of nanomole amounts by 600-mhz (1)h and (13)c NMR analysis. *J Biol Chem* 275: 3896-3906.
- Guerry P (1997) Nonlipopolysaccharide surface antigens of *Campylobacter* species. *J Infect Dis* 176: S122-124.
- Hamali H, Fallah S, Joozani RJ, Zare P, Noorsaadat G (2014) Detection of *Campylobacter* spp. in sheep aborted fetuses by PCR. *Trends Life Sci* 3: 49-56.
- Harvey SM, Greenwood JR (1983) Relationships among catalase-positive *campylobacters* determined by deoxyribonu-

- cleic acid-deoxyribonucleic acid hybridization. *Int J Syst Evol Microbiol* 33: 275-284.
- Hedstrom OR, Sonn RJ, Lassen ED, Hultgren BD, Crisman RO, Smith BB, Snyder SP (1987) Pathology of *Campylobacter jejuni* abortion in sheep. *Vet Pathol* 24: 419-426.
- Hickey TE, McVeigh AL, Scott DA, Michielutti RE, Bixby A, Carroll SA, Bourgeois AL, Guerry P (2000) *Campylobacter jejuni* cytolethal distending toxin mediates release of interleukin-8 from intestinal epithelial cells. *Infect Immun* 68: 6535-6541.
- Jeon B, Itoh K, Ryu S (2005) Promoter analysis of cytolethal distending toxin genes (*cdtA*, B, and C) and effect of a *luxS* mutation on CDT production in *Campylobacter jejuni*. *Microbiol Immunol* 49: 599-603.
- Joens LA, Haesebrouck F, Pasmans F (2010) *Campylobacter* and *Helicobacter* In: Gyles CL, Prescott JF, Songer JG, Theon CO (Eds). *Pathogenesis of Bacterial Infections in Animals*. *Med Microbiol* (4th edition pp. 483-501). Wiley-Blackwell Publishing
- Kirkbride CA (1993) Diagnoses in 1784 ovine abortions and stillbirths. *J Vet Diagn Invest* 5: 398-402.
- Konkel ME, Gray SA, Kim BJ, Garvis SG, Yoon J (1999) Identification of the enteropathogens *Campylobacter jejuni* and *Campylobacter coli* based on the *cadF* virulence gene and its product. *J Clin Microbiol* 37: 510-517.
- McFadyean J, Stockman S (1913) Report of the departmental committee appointed by the board of agriculture and fisheries to enquire into epizootic abortion, part iii (pp 1-64). Her Majesty's Stationary office, London. www.google.co.in/books/edition/Report_of_the_Departmental_Committee.
- Mearns R (2007) Abortion in sheep 1. Investigation and principal causes. *In Practice* 29: 40-46.
- Monteville MR, Yoon JE, Konkel ME (2003) Maximal adherence and invasion of INT 407 cells by *Campylobacter jejuni* requires the CadF outer-membrane protein and microfilament reorganization. *Microbiol* 149: 153-165.
- Muller J, Schulze F, Muller W, Hänel I (2006) PCR detection of virulence-associated genes in *Campylobacter jejuni* strains with differential ability to invade Caco-2 cells and to colonize the chick gut. *Vet Microbiol* 113: 123-129.
- Padungton P, Kaneene JB (2003) *Campylobacter* spp. in human, chickens, pigs and their antimicrobial resistance. *J Vet Med Sci* 65: 161-70.
- Perez-Perez, GI, Blaser, MJ (1996) *Campylobacter* and *Helicobacter*. Carlton LG, John F, Prescott JF, Songer JG, Thoen CO, (Eds). *Pathogenesis of Bacterial Infections in Animals*. *Med Microbiol* (4th edition). pp. 483-501. Blackwell Publishing
- Poropatich KO, Walker CLF, Black RE (2010) Quantifying the association between *Campylobacter* infection and Guillain-Barré syndrome: a systematic review. *J Health Popul Nutr* 28: 545-552
- Rahimi E, Kazemini HR, Safaei S, Allahbakhshi K, Momeni M, Riahi M (2010) Detection and identification of *Campylobacter* spp. from retail raw chicken, turkey, sheep and goat meat in Ahvaz, Iran. *Afr J Microbiol Res* 4: 1620-1623.
- Raji MA, Adekeye JO, Kwaga JK, Bale JO (2000) Bioserogroups of *Campylobacter* species isolated from sheep in Kaduna State, Nigeria. *Small Rumin Res* 37: 215-221.
- Reay DJM (1908) Report of the Departmental Committee appointed by the Board of Agriculture and Fisheries to inquire into and report upon the subject of agricultural education in England and Wales. In: Parliament., Printed for HMSO, by McCorquodale & Co., Ltd.. 1908.
- Reddy S, Zishiri OT (2018) Genetic characterisation of virulence genes associated with adherence, invasion and cytotoxicity in *Campylobacter* spp. isolated from commercial chickens and human clinical cases. *Onderstepoort J Vet Res* 85: 1507.
- Rozynek E, Dzierzanowska-Fangrat K, Jozwiak P, Popowski J, Korsak D, Dzierzanowska D (2005) Prevalence of potential virulence markers in Polish *Campylobacter jejuni* and *Campylobacter coli* isolates obtained from hospitalized children and from chicken carcasses. *J Med Microbiol* 54: 615-619.
- Sahin O, Plummer PJ, Jordan DM, Sulaj K, Pereira S, Robbe-Austerman S, Wang L, Yaeger MJ, Hoffman LJ, Zhang Q (2008) Emergence of a tetracycline-resistant *Campylobacter jejuni* clone associated with outbreaks of ovine abortion in the United States. *J Clin Microbiol* 46: 1663-1671.
- Sahin O, Yaeger M, Wu Z, Zhang Q (2017) *Campylobacter*-associated diseases in animals. *Annu Rev Anim Biosci* 5: 21-42.
- Salama SM, Garcia MM, Taylor DE (1992) Differentiation of the subspecies of *Campylobacter fetus* by genomic sizing. *Int J Syst Evol Microbiol* 42: 446-450.
- Steinhauserova I, Ceskova J, Fojtikova K, Obrovskaa I (2001) Identification of thermophilic *Campylobacter* spp. by phenotypic and molecular methods. *J Appl Microbiol* 90: 470-475.
- Wang G, Clark CG, Taylor TM, Pucknell C, Barton C, Price L, Woodward DL, Rodgers FG (2002) Colony multiplex PCR assay for identification and differentiation of *Campylobacter jejuni*, *C. coli*, *C. lari*, *C. upsaliensis*, and *C. fetus* subsp. *fetus*. *J Clin Microbiol* 40: 4744-4747.
- Wieczorek K, Osek J (2013) Antimicrobial resistance mechanisms among *Campylobacter*. *Bio Med Res Int* doi: 10.1155/2013/340605.
- Wieczorek K, Osek J (2008) Identification of virulence genes in *Campylobacter jejuni* and *C. coli* isolates by PCR. *Bull Vet Inst Pulawy* 52: 211-216.
- Xue F, Jiang Y, Xu F, Zhang X, Luan J, Zhu C, Chen H (2009) Isolation and characterization of *Campylobacter* from red-crowned cranes in China. *J Anim Vet Adv* 8: 2442-2446.
- Yaeger MJ, Sahin O, Plummer PJ, Wu Z, Stasko JA, Zhang Q (2021) The pathology of natural and experimentally induced *Campylobacter jejuni* abortion in sheep. *J Vet Diagn Invest* 33: 1096-1105.
- Yoo JH, Choi NY, Bae YM, Lee JS, Lee SY (2014) Development of a selective agar plate for the detection of *Campylobacter* spp. in fresh produce. *Int J Food Microbiol* 189: 67-74.
- Yuki N (2001) Infectious origins of, and molecular mimicry in, Guillain-Barré and Fisher syndromes. *Lancet Infect Dis* 1: 29-37.
- Zilbauer M, Dorrell N, Wren BW, Bajaj-Elliott M (2008) *Campylobacter jejuni*-mediated disease pathogenesis: an update. *Trans R Soc Trop Med Hyg* 102: 123-129.
- Ziprin RL, Young CR, Byrd JA, Stanker LH, Hume ME, Gray SA, Kim BJ, Konkel ME (2001) Role of *Campylobacter jejuni* potential virulence genes in cecal colonization. *Avian Dis* 45: 549-557.